



## Guideline Summary NGC-6536

### Guideline Title

**Treatment of urinary tract infections in nonpregnant women.**

### Bibliographic Source(s)

American College of Obstetricians and Gynecologists (ACOG). Treatment of urinary tract infections in nonpregnant women. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 2008 Mar. 10 p. (ACOG practice bulletin; no. 91). [51 references]

### Guideline Status

This is the current release of the guideline.

### FDA Warning/Regulatory Alert

**Note from the National Guideline Clearinghouse:** This guideline references a drug(s) for which important revised regulatory and/or warning information has been released.

- [August 15, 2013 - Fluoroquinolone Antibacterial Drugs](#) : The U.S. Food and Drug Administration (FDA) has required the drug labels and Medication Guides for all fluoroquinolone antibacterial drugs be updated to better describe the serious side effect of peripheral neuropathy. This serious nerve damage potentially caused by fluoroquinolones may occur soon after these drugs are taken and may be permanent.

### Scope

#### Disease/Condition(s)

- Urinary tract infections
- Acute bacterial cystitis
  - Acute pyelonephritis

#### Guideline Category

- Diagnosis  
Evaluation  
Management  
Prevention  
Risk Assessment  
Screening  
Treatment

#### Clinical Specialty

- Family Practice  
Internal Medicine  
Obstetrics and Gynecology  
Urology

#### Intended Users

- Physicians

#### Guideline Objective(s)

- To aid practitioners in making decisions about appropriate obstetric and gynecologic care
- To address the diagnosis, treatment, and prevention of uncomplicated acute bacterial cystitis and acute bacterial

pyelonephritis in nonpregnant women

## Target Population

Nonpregnant women with uncomplicated urinary tract infections (UTIs)

**Note:** Women with complicated UTIs (e.g., in patients with diabetes mellitus, abnormal anatomy, prior urologic surgery, a history of renal stones, an indwelling catheter, spinal cord injury, immunocompromise, or in pregnant patients) are a heterogeneous group of conditions beyond the scope of this guideline.

## Interventions and Practices Considered

### Evaluation and Diagnosis

1. Differential diagnosis: acute bacterial cystitis versus acute pyelonephritis
  - Clinical history and physical examination
  - Laboratory evaluation of bacteriuria and pyuria
    - Urine dipstick testing
    - Urine culture or urinalysis
2. Risk factor assessment for urinary tract infection (UTI) in premenopausal and postmenopausal women
3. Risk factor assessment for recurrent UTI
4. Imaging of the urinary tract (not recommended routinely for uncomplicated UTIs)
5. Screening for asymptomatic bacteriuria in nonpregnant, premenopausal women (considered but specifically not recommended)

### Treatment and Management

1. Antimicrobial regimens for uncomplicated UTIs (3-day vs. 7-day)
  - Trimethoprim-sulfamethoxazole
  - Trimethoprim
  - Ciprofloxacin
  - Levofloxacin
  - Norfloxacin
  - Gatifloxacin
  - Nitrofurantoin macrocrystals and nitrofurantoin monohydrate macrocrystals
  - Fosfomycin tromethamine
2. Use of 14-day antimicrobial regimens for acute pyelonephritis
3. Management of recurrent UTI
  - Prophylactic or intermittent antimicrobial therapy
  - Cranberry juice
  - Methenamine salts (not sufficient evidence for use)
4. Treatment of UTIs in postmenopausal women

## Major Outcomes Considered

- Incidence of acute bacterial cystitis
- Incidence of acute pyelonephritis
- Incidence of recurrent urinary tract infections
- Effectiveness of antimicrobial therapy (clinical response and bacterial eradication rates)
- Sensitivity and specificity of diagnostic tests
- Adverse effects of treatment

## Methodology

### Methods Used to Collect/Select the Evidence

Hand-searches of Published Literature (Primary Sources)

Hand-searches of Published Literature (Secondary Sources)

Searches of Electronic Databases

### Description of Methods Used to Collect/Select the Evidence

The MEDLINE database, the Cochrane Library, and the American College of Obstetricians and Gynecologists' own internal resources and documents were used to conduct a literature search to locate relevant articles published between January 1995 and April 2007. The search was restricted to articles published in the English language. Restrictions were placed on articles

1985 and April 2007. The search was restricted to articles published in the English language. Priority was given to articles reporting results of original research, although review articles and commentaries also were consulted. Abstracts of research presented at symposia and scientific conferences were not considered adequate for inclusion in this document. Guidelines published by organizations or institutions such as the National Institutes of Health and the American College of Obstetricians and Gynecologists were reviewed, and additional studies were located by reviewing bibliographies of identified articles.

### Number of Source Documents

Not stated

### Methods Used to Assess the Quality and Strength of the Evidence

Weighting According to a Rating Scheme (Scheme Given)

### Rating Scheme for the Strength of the Evidence

Studies were reviewed and evaluated for quality according to the method outlined by the U.S. Preventive Services Task Force:

**I** Evidence obtained from at least one properly designed randomized controlled trial.

**II-1** Evidence obtained from well-designed controlled trials without randomization.

**II-2** Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.

**II-3** Evidence obtained from multiple time series with or without the intervention. Dramatic results in uncontrolled experiments also could be regarded as this type of evidence.

**III** Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

### Methods Used to Analyze the Evidence

Review of Published Meta-Analyses

Systematic Review

### Description of the Methods Used to Analyze the Evidence

Not stated

### Methods Used to Formulate the Recommendations

Expert Consensus

### Description of Methods Used to Formulate the Recommendations

Analysis of available evidence was given priority in formulating recommendations. When reliable research was not available, expert opinions from obstetrician-gynecologists were used. See also the "Rating Scheme for the Strength of Recommendations" field regarding Grade C recommendations.

### Rating Scheme for the Strength of the Recommendations

Based on the highest level of evidence found in the data, recommendations are provided and graded according to the following categories:

**Level A** — Recommendations are based on good and consistent scientific evidence.

**Level B** — Recommendations are based on limited or inconsistent scientific evidence.

**Level C** — Recommendations are based primarily on consensus and expert opinion.

### Cost Analysis

Published cost analyses were reviewed.

### Method of Guideline Validation

Internal Peer Review

### Description of Method of Guideline Validation

Practice Bulletins are validated by two internal clinical review panels composed of practicing obstetrician-gynecologists generalists and sub-specialists. The final guidelines are also reviewed and approved by the American College of Obstetricians and Gynecologists (ACOG) Executive Board.

## Recommendations

### Major Recommendations

The grades of evidence (I-III) and levels of recommendation (A-C) are defined at the end of the "Major Recommendations" field.

**The following recommendations and conclusions are based on good and consistent scientific evidence (Level A):**

- Screening for and treatment of asymptomatic bacteriuria is not recommended in nonpregnant, premenopausal women.
- Resistance rates higher than 15 to 20% necessitate a change in antibiotic class.
- In all cases of acute pyelonephritis, whether treatment is on an inpatient or outpatient basis, 14 days of total antimicrobial therapy should be completed.
- A 3-day antimicrobial regimen is the preferred treatment duration for uncomplicated acute bacterial cystitis in women, including women aged 65 years and older.

**The following conclusion is based on limited or inconsistent evidence (Level B):**

- The initial treatment of a symptomatic lower urinary tract infection (UTI) with pyuria or bacteriuria or both does not require a urine culture.

**The following conclusions are based primarily on consensus and expert opinion (Level C):**

- Beta-lactams, such as first-generation cephalosporins and amoxicillin, are less effective in the treatment of acute uncomplicated cystitis than those antimicrobials listed in the Table below.
- To diagnose bacteriuria, decreasing the colony count to 1,000 to 10,000 bacteria per milliliter in symptomatic patients will improve the sensitivity without significantly compromising specificity.

**Table. Treatment Regimens for Uncomplicated Acute Bacterial Cystitis**

Antimicrobial Agent	Dose	Adverse Events
Trimethoprim-sulfamethoxazole	One tablet (160 mg trimethoprim-800 mg sulfamethoxazole), twice daily for 3 days	Fever, rash, photosensitivity, neutropenia, thrombocytopenia, anorexia, nausea and vomiting, pruritus, headache, urticaria, Stevens-Johnson syndrome, and toxic epidermal necrosis
Trimethoprim	100 mg, twice daily for 3 days	Rash, pruritus, photosensitivity, exfoliative dermatitis, Stevens-Johnson syndrome, toxic epidermal necrosis, and aseptic meningitis
Ciprofloxacin	250 mg, twice daily for 3 days	Rash, confusion, seizures, restlessness, headache, severe hypersensitivity, hypoglycemia, hyperglycemia, and Achilles tendon rupture (in patients older than 60 years)
Levofloxacin	250 mg, once daily for 3 days	Same as for ciprofloxacin
Norfloxacin	400 mg, twice daily for 3 days	Same as for ciprofloxacin
Gatifloxacin	200 mg, once daily for 3 days	Same as for ciprofloxacin
Nitrofurantoin macrocrystals	50 to 100 mg, four times daily for 7 days	Anorexia, nausea, vomiting, hypersensitivity, peripheral neuropathy, hepatitis, hemolytic anemia, and pulmonary reactions
Nitrofurantoin monohydrate crystals	100 mg, twice daily for 7 days	Same as for nitrofurantoin macrocrystals
Fosfomycin tromethamine	3 g dose (powder) single dose	Diarrhea, nausea, vomiting, rash, and hypersensitivity

**Definitions:**

**Grades of Evidence**

**I** Evidence obtained from at least one properly designed randomized controlled trial.

**II-1** Evidence obtained from well-designed controlled trials without randomization.

**II-2** Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.

**II-3** Evidence obtained from multiple time series with or without the intervention. Dramatic results in uncontrolled experiments also could be regarded as this type of evidence.

**III** Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

**Levels of Recommendation**

**Level A** — Recommendations are based on good and consistent scientific evidence.

**Level B** — Recommendations are based on limited or inconsistent scientific evidence.

**Level C** — Recommendations are based primarily on consensus and expert opinion.

**Clinical Algorithm(s)**

None provided

**Evidence Supporting the Recommendations**

**Type of Evidence Supporting the Recommendations**

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

**Benefits/Harms of Implementing the Guideline Recommendations**

**Potential Benefits**

Accurate diagnosis and appropriate management of urinary tract infections in nonpregnant women

### Potential Harms

Adverse events associated with antimicrobial treatment regimens (see Table 1 in the original guideline document).

### Qualifying Statements

#### Qualifying Statements

These guidelines should not be construed as dictating an exclusive course of treatment or procedure. Variations in practice may be warranted based on the needs of the individual patient, resources, and limitations unique to the institution or type of practice.

### Implementation of the Guideline

#### Description of Implementation Strategy

An implementation strategy was not provided.

#### Implementation Tools

Audit Criteria/Indicators

Foreign Language Translations

Patient Resources

For information about availability, see the *Availability of Companion Documents* and *Patient Resources* fields below.

### Institute of Medicine (IOM) National Healthcare Quality Report Categories

#### IOM Care Need

Getting Better

Staying Healthy

#### IOM Domain

Effectiveness

### Identifying Information and Availability

#### Bibliographic Source(s)

American College of Obstetricians and Gynecologists (ACOG). Treatment of urinary tract infections in nonpregnant women. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 2008 Mar. 10 p. (ACOG practice bulletin; no. 91). [51 references]

#### Adaptation

Not applicable: The guideline was not adapted from another source.

#### Date Released

2008 Mar

#### Guideline Developer(s)

American College of Obstetricians and Gynecologists - Medical Specialty Society

#### Source(s) of Funding

American College of Obstetricians and Gynecologists (ACOG)

#### Guideline Committee

American College of Obstetricians and Gynecologists (ACOG) Committee on Practice Bulletins - Gynecology

#### Composition of Group That Authored the Guideline

Not stated

#### Financial Disclosures/Conflicts of Interest

Not stated

## Guideline Status

This is the current release of the guideline.

## Guideline Availability

Electronic copies: None available

Print copies: Available for purchase from the American College of Obstetricians and Gynecologists (ACOG) Distribution Center, PO Box 933104, Atlanta, GA 31193-3104; telephone, 800-762-2264, ext. 192; e-mail: [sales@acog.org](mailto:sales@acog.org). The ACOG Bookstore is available online at the [ACOG Web site](#).

## Availability of Companion Documents

Proposed performance measures are included in the original guideline document.

## Patient Resources

The following is available:

- Urinary tract infections. Atlanta (GA): American College of Obstetricians and Gynecologists (ACOG); 2006.

Electronic copies: Available from the [American College of Obstetricians and Gynecologists \(ACOG\) Web site](#). Copies are also available in Spanish.

Print copies: Available for purchase from the American College of Obstetricians and Gynecologists (ACOG) Distribution Center, PO Box 933104, Atlanta, GA 31193-3104; telephone, 800-762-2264, ext. 192; e-mail: [sales@acog.org](mailto:sales@acog.org). The ACOG Bookstore is available online at the [ACOG Web site](#).

Please note: This patient information is intended to provide health professionals with information to share with their patients to help them better understand their health and their diagnosed disorders. By providing access to this patient information, it is not the intention of NGC to provide specific medical advice for particular patients. Rather we urge patients and their representatives to review this material and then to consult with a licensed health professional for evaluation of treatment options suitable for them as well as for diagnosis and answers to their personal medical questions. This patient information has been derived and prepared from a guideline for health care professionals included on NGC by the authors or publishers of that original guideline. The patient information is not reviewed by NGC to establish whether or not it accurately reflects the original guideline's content.

## NGC Status

This NGC summary was completed by ECRI Institute on July 30, 2008. The information was verified by the guideline developer on August 20, 2008. This summary was updated by ECRI Institute on October 25, 2013 following the U.S. Food and Drug Administration advisory on Fluoroquinolone Antibacterial Drugs.

## Copyright Statement

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

## Disclaimer

### NGC Disclaimer

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion-criteria.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.