

MONTHLY CODING SPOTLIGHT



MAJOR DEPRESSIVE DISORDER

In ICD-10, Major Depressive Disorder is categorized by episode and/or severity.

MDD either occurs in a single episode or it is recurrent. These episodes can be further defined by severity of mild, moderate or severe. PHQ-9 is a scoring tool designed to help assess these levels of severity.

In severe depression, psychotic features may be detectable and should be reported to that higher level of specificity.

When MDD is controlled it doesn't mean it has gone away. It can still be diagnosed in a remission status, either fully or partially.

Bipolar disorder can be reported similarly to MDD with mild, moderate or severe episodes, including psychotic features in the severe cases and in partial or full remission.

Please note that Bipolar disorder should not be coded with Major Depression.

Schizophrenia is also a significant mental disorder and is reported by the type of schizophrenia: paranoid, disorganized, catatonic, etc.

Example:

A/P: Mild episode of recurrent major depressive disorder. This is stable, continue Mirtazapine tablet, once by mouth every night at bedtime.

Diagnosis:

⇒ F33.0 Major depressive disorder, recurrent, mild

With all mental disorders, be sure to document any dependency on medications, cannabis or alcohol.

AZPC Coding and Documentation materials are based on current guidelines and are to be used for reference only. Clinical and coding decisions are to be made based on the independent judgement of the treating physician or qualified health care practitioner and the best interests of the patient. ICD-10-CM, CPT and HCPCS are the authoritative references for purposes of assigning diagnoses and procedure codes to be reported. It is the responsibility of the physician and/or coding staff to determine and submit accurate codes, charges and modifiers for services rendered.

MDD CODES AT A GLANCE



- F20._ Schizophrenia
- F25._ Schizoaffective Disorders
- F30. Manic Episodes
- F31. Bipolar Disorders
- F32. Major Depressive Disorder, single episode
 - F32.0 MDD, single episode, mild
 - F32.1 MDD, single episode, moderate
 - F32.2 MDD, single episode, severe without psychotic features
 - F32.3 MDD, single episode, severe with psychotic features
 - F32.4 MDD, single episode, in partial remission
 - F32.5 MDD, single episode, in full remission
 - F32.9 MDD, unspecified (not a preferred code)

F33. Major Depressive Disorder, recurrent

- F33.0 MDD, recurrent, mild
- F33.1 MDD, recurrent, moderate
- F33.2 MDD, recurrent, severe without psychotic features
- F33.3 MDD, recurrent, severe with psychotic features
- F33.4_ Major Depressive Disorder, recurrent, in remission
 - F33.40 MDD, recurrent, in remission unspecified
 - F33.41 MDD, recurrent, in partial remission
 - F33.42 MDD, recurrent, in full remission
 - F33.8 Other recurrent depressive disorders
 - F33.9 MDD, recurrent, unspecified

F34.8 Other Persistent Mood (affective) Disorders

F34.9 Unspecified Mood (affective) Disorder

- F10.20 Alcohol Dependence Uncomplicated
- F11.20 Opioid Dependence Uncomplicated
- F12.20 Cannabis Dependence Uncomplicated

AZPC Coding and Documentation materials are based on current guidelines and are to be used for reference only. Clinical and coding decisions are to be made based on the independent judgement of the treating physician or qualified health care practitioner and the best interests of the patient. ICD-10-CM, CPT and HCPCS are the authoritative references for purposes of assigning diagnoses and procedure codes to be reported. It is the responsibility of the physician and/or coding staff to determine and submit accurate codes, charges and modifiers for services rendered.